PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:
C12N 15/85, C12Q 1/68

A1

(11) International Publication Number: WO 99/66060

(43) International Publication Date: 23 December 1999 (23.12.99)

(21) International Application Number: PCT/IB99/01071

(22) International Filing Date: 11 June 1999 (11.06.99)

(30) Priority Data:

10/170941

18 June 1998 (18.06.98)

JР

(71) Applicant (for all designated States except US): HOECHST MARION ROUSSEL LTD. [JP/JP]; 17-51, Akasaka 2-chome, Minato-ku, Tokyo 107-8465 (JP).

(72) Inventors: and


- (75) Inventors/Applicants (for US only): KAWAI, Shinji [JP/FR]; 416, rue Emile Dubois, F-75014 Paris (FR). SUGIURA, Takeyuki [JP/JP]; 4-4-22-202, Kamishakuji-i, Nerima-ku, Tokyo 177-0044 (JP). HÖTTEN, Gertrud [DE/DE]; Hölkeskampring 96, D-44625 Herne (DE).
- (74) Agent: VIEILLEFOSSE, Jean-Claude; Hoechst Marion Roussel, 102, route de Noisy, F-93235 Romainville Cedex (FR).

(81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: HUMAN MP52 GENE PROMOTER AND METHOD FOR EXPLORING USEFUL SUBSTANCE BY USING THE SAME



OKST AVAILABLE COR,

(57) Abstract

The present object is to provide a method for exploring low molecular weight compounds having osteogenetic activity or an activity to inhibit osteogenesis that shows an equal degree to human BMP proteins. A method for exploring low molecular weight compounds which regulate positively or negatively the expression of human MP52 with reference to a reporter activity by using 5' upstream region gene containing the human MP52 promoter and an animal cell into which a recombinant expression vector that has been ligated to an appropriate reporter gene has been introduced. Low molecular weight compounds and their derivatives obtained by the present method have morphogenetic activity and inhibiting activity for bone and cartilage, inducing or inhibiting activity for tendon and ligament, forming, inducing, and protecting effects on neural cells, or inducing or inhibiting angiogenic activity, and are effective as preventive or therapeutic agent for cartilage and bone diseases, therapeutic agent for injury of tendon and ligament, therapeutic agent for neural diseases, or healing agent for injury and carcinostatic, through the expression of human MP52.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	· TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali .	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukrainė
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Сапада	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania .		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		•
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

1

HUMAN MP52 GENE PROMOTER AND METHOD FOR EXPLORING USEFUL SUBSTANCE BY USING THE SAME

BACKGROUND OF THE INVENTION

5 (1) Field of the Invention

The present invention relates to a 5' upstream region DNA sequence which contains a promoter of human MP52 gene. Furthermore, the present invention provides a method for exploring low molecular weight compound regulating positively or negatively the expression of human MP52 by using a mass of animal cells transfected with a recombinant expression vector harboring the 5' upstream region DNA containing the human MP52 gene promoter in front of a suitable reporter gene, and by using a reporter activity as an indicator.

15 (2) Description of the Related Art

Many substances have been known as usable for healing of bone deficit or fracture, by enhancing bone regeneration.

Among them, a bone morphogenetic protein (BMP, hereafter) is a protein belonging to TGF (transforming growth factor)

 $-\beta$ superfamily and has been formerly found as a substance inducing ectopic ossification and existing in decalcified bone tissue (Trends in Biotechnol. 11, 379-383, 1993). BMP members known so far range from BMP-1 to BMP-14. Among them, the members from BMP-2 to BMP-14 have been known as showing the bone morphogenetic activity.

Human MP52 is a BMP-like protein isolated by PCR cloning (Biochem. Biophys. Res. Comm. 204, 646-652, 1994).

Recombinant human MP52 has been reported as effective not only in inducing ectopic ossification, but also in curing bone deficit in an animal model (Growth Factors 13, 65-74, 1996).

It is considered that these BMP proteins are effective in treating and preventing various bone disorders and bone diseases. However, the BMP proteins exist in very small quantity in nature, and for an available large quantity for therapeutic use, manufacturing a recombinant protein is necessary.

Preparing a recombinant protein is generally very

PCT/IB99/01071

WO:99/66060

20

expensive compared with low molecular weight compounds. Furthermore, there are many restrictions as a medicinal drug in terms of physical properties and administration methods due to proteineous characteristics. Considering these 5 points, a small molecular organic compound having the activity equal to that of the BMP proteins described above, if any, should be a highly promising medicinal drug. SUMMARY OF THE INVENTION

The present invention provides a method for exploring a 10 substance which induces human MP52, a BMP-like protein. useful substance obtained by this exploring method can be effective in inducing expression of human MP52, a bone formation factor. Since it has a small molecular weight and the same activity as that of human MP52, it can have a very 15 useful application. On the contrary, the present invention provides a method for exploring a substance which inhibits expression of human MP52. In case that human MP52 relates to bone and cartilage hyperplasia, it can prevent hyperplasia by inhibiting the expression of human MP52.

Recently, it has been suggested that a mutant of the human MP52 protein not only decreases the activity per se, but also deactivates other BMPs by forming dimer proteins (Nature Genetics, 17, 58-64, 1997). The substance inhibiting the expression of human MP52 may be effective in mitigating 25 chondrodystrophy caused by such a mutant MP52.

Moreover, it has been known that human MP52 has a tendon and ligament inducing activity (Toshiyuki Tashiro et al., American Society of Bone and Mineral Research, F336, 1997), neural cells forming, inducing, and protecting effects (J. 30 Neurosci. Res., 42, 724-732, 1995), and angiogenic activity (Exp. Cell Res., 235, 218-226, 1997). Thus, it is possible that the exploring method provided by the present invention gives tendon and ligament inducing and inhibiting substances, neural inducing substances, or pro- and anti-angiogenic 35 substances.

For such an exploring method, an example was so far only reported using a mouse BMP-2 promoter (W097/15308); there is no example of using the human MP52 promoter. In addition,

since the materials of the exploring method provided by the present invention are all derived from human sources, it can be expected that the discovered compounds should show the clinical effects.

5 BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a 3.5 kb of 5' upstream region of human MP52 gene and a restriction enzyme map.

Fig. 2 is a recombinant expression vector (pMP52L) containing a 5' upstream region of human MP52 gene. A 10 fragment, base No. from 1 to 3227 (Apal position 1) shown in SEQ ID No.: 1 of the Sequence Listing, was inserted to the restriction enzyme site, HindIII-KpnI, of pGL3-basic.

Fig. 3 is a result of measuring a human MP52 promoter activity (transient expression). pGL3B means pGL3-basic in the figure.

DESCRIPTION OF THE PREFERRED EMBODIMENT

The present invention relates to a DNA whose nucleotide sequence is represented by the base sequence from No. 1 to No. 3521 in SEQ ID NO.: 1 of the Sequence Listing that 20 encodes a human MP52 gene promoter region, or a fragment thereof. SEQ ID NO.: 1 of the Sequence Listing shows a 5' upstream region sequence of the human MP52 gene.

The present invention relates to a method for preparing the DNA shown in SEQ ID NO.: 1 of the Sequence Listing having 25 the steps of:

- (1) preparation of a cDNA probe by PCR with a degenerate primer,
- (2) plaque hybridization screening of a human genomic phage library by using the probe, and
- 30 (3) subcloning of a target fragment from an isolated phage clone.

The plasmid vector used here is not restricted and any commercialized vector can be used. A preferable example is pUC18.

35 The present invention relates to a recombinant expression vector characterized by integration of the full length or a part of DNA shown in SEQ ID NO.: 1 of the Sequence Listing into a reporter gene. In detail, the

4

recombinant expression vector is constructed to locate a suitable region of 5' upstream region of the human MP-52 gene, that is represented by SEQ ID No.: 1 of the Sequence Listing, in front of a reporter gene. The reporter gene such 5 as luciferase or β -galactosidase gene shows an expressing status on behalf of an original product. The basic vector for constructing the recombination expression vector is not specially restricted to allow to use a plasmid vector The present invention uses pGL3-basic as a commercialized. 10 preferable example. The use of pGL3-basic yielded a pMP52L (8085 bp) that is a recombination vector containing the human MP52 promoter and a luciferase reporter gene. The present invention assigned it to a recombination expression vector. It is necessary to introduce the vector, to mammalian cells, 15 preferably a human osteoblast-like cells, such as SaOS-2 cells, with a liposome. The animal cells stably transfected with the recombinant expression vector are selected by using a resistance marker.

The present invention relates to an exploring method for 20 a useful substance by using a recombinant expression vector characterized by integrating the full length or a part of DNA shown in SEQ ID NO .: 1 of the Sequence Listing into a reporter gene. The sequence from 2880 to 3521 in SEQ ID NO.: 1 of the Sequence Listing has been already reported (Biochem. 25 Biophys. Res. Comm., 204, 646-652, 1994). The longer sequence has been integrated in a recombinant expression vector, the more natural process of the exploring method can be used. The present invention relates to the method for exploring a useful substance which is a bone-related 30 substance. In particular, the present invention relates to the method for exploring a useful substance which is an osteogenesis inducing substance or which is an osteogenesis inhibiting substance. Furthermore, the present invention relates to the method for exploring a useful substance which 35 is a neurogenic substance or which is an angiogenic substance. A low molecular weight compound which induces or inhibits the expression of human MP52 can be obtained by isolating the promoter which regulates the expression of the

gene, by ligating it to a suitable reporter gene and by introducing the gene structure into a suitable mammal cell to make an exploring system. The substance which regulates the expression of human MP52 in the exploring system works on the promoter to increase or decrease the expression level of the reporter gene. Therefore, a simple and easy measurement of the reporter activity makes an exploration of the aimed possible substance.

The animal cell transfected with said vector can be used 10 for a method for screening a chemical compound library by high throughput screening (Nature, 384, supp, 14-16, 1996) and finding an active substance from natural compounds. The substance which increases or decreases an activity is searched by treating the cell with a compound for an 15 appropriate period of time and measuring a reporter activity. The compound obtained in such a way can regulate the expression by working directly on a transcription factor or indirectly on the promoter of human MP52 through regulating a signal transduction system. Therefore, these compounds are 20 effective as a therapeutic agent for osteocartilaginous diseases, cancer metastasis to bone, or osteohyperplasia. other examples, these compounds are effective as a recovering agent for tendon and ligament injuries, a remedy for neural diseases, or an angiogenesis inducing or inhibiting agent.

25 The substance obtained by the present invention has a bone or cartilage morphogenetic activity and is effective as an agent for therapeutic and preventive treatment in the field of orthopedic surgery (fracture, osteoarthritis such as joint osteoarthritis and hip joint osteoarthritis, at arthrosteitis, damage of cartilage such as damage of meniscus, regeneration of bone and cartilage deficit caused by injury and tumor dissection, bone reconstruction such as spinal fusion and vertebral canal enlargement, and congenital cartilage and bone diseases such as dysosteogenesis and achondroplasia), or dental fields (bone reconstruction such as palatoschisis, mandible reconstruction, and residual ridge construction), and osteoporosis. Moreover, the substance of the present invention can be used for bone graft in aesthetic

surgery. These therapeutic treatments are effective in treating the field of veterinary surgery. On the other hand, the present invention can provide a substance which inhibits bone or cartilage morphogenesis. In this case, the substance 5 is applied as an agent to prevention and therapy of bone or cartilage hyperplasia.

Moreover, the substance which induces or inhibits tendon and ligament can be obtained according to the present invention. Therefore, the present invention provides a 10 method for exploring medicinal drugs to treat damage of tendons and ligaments. Furthermore, since human MP52 has neural cells forming, inducing, and protecting effects, the compounds obtained from the present invention can be applied as therapeutic agents for neural diseases. In addition, it 15 is possible to obtain a substance inducing or inhibiting angiogenesis. The compounds obtained can be used as remedy for injury, carcinostatic and metastasis inhibitor.

By using 5' upstream region containing the human MP52 gene promoter and an animal cell into which a recombinant 20 expression vector that has been integrated in an appropriate reporter gene, low molecular weight compounds regulating positively or negatively the expression of human MP52 can be explored with reference to a reporter activity. molecular weight compounds and their derivatives have 25 morphogenetic activity and inhibiting activity for bone and cartilage through the expression of human MP52 and are useful as preventive or therapeutic agents for cartilage and bone diseases, remedies for osteometastasis, or therapeutic and preventive agents for osteohyperplasia. Furthermore, these 30 compounds are effective as repairing agent for injury of tendon and ligament, therapeutic agent for neural diseases, and inducing and inhibiting agent for angiogenesis.

EXAMPLES

This invention shall be more illustratively explained by 35 way of the following Examples. The following Examples are to be considered in all respects as illustrative and not restrictive.

Example 1 : Cloning of 5' upstream region of human MP52 gene

7

The 40 cycles of PCR reaction was carried out by using the primer OD of SEQ ID No.: 2 of the Sequence Listing and the primer OID of SEQ ID No.: 3 of the Sequence Listing, cDNA derived from a human fetus (8-9 week old) as a template, and 5 1.5 unit Taq polymerase (provided from Perkin Elmer Cetus). The following processes were repeated twice: the DNA amplified was treated with SphI and AlwNI and then the PCR reaction was carried out 13 times. Finally, it was treated with EcoRI, the PCR product was ligated to pBluescript SK 10 (provided from Stratagene Ltd.). A human genomic phage library (provided from Stratagene Ltd.) was screened by plaque hybridization using the cDNA sequence of MP52 among cloned sequences. A 20 kb DNA obtained was subcloned to pBluescript SK vector, and an unnecessary part to remove was 15 cleaved by restriction with SalI and BamHI, to obtain the pBluescript vector containing a human MP52 DNA of 8.8 kb. The vector was named E. coli pMP52. E. coli pMP52 was deposited in National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, 20 Ministry of International Trade and Industry 1-3, Higashi 1chome, Tsukuba-shi Ibaraki-ken 305-8566 Japan, in March 30, 1998 with depository No. FERM P-16734 and transferred to the International Depository Authority under Budapest Treaty on April 1, 1999 (Deposit No. FERM BP-6688).

25 <u>Example 2</u>: Determination of DNA sequence of 5' upstream region of human MP52 gene

The pBluescript vector containing a human MP52 DNA of 8.8 kb was digested by XbaI, a remained large fragment (7.6 kb) was purified by agarose electrophoresis to subject to self-ligation by using a ligation kit provided from Takara Shuzo Ltd. The 3.5 kb sequence of the 5' terminal was determined by using a vector containing the upstream region of the human MP52 gene of 4.7 kb as a template and by a ALF automated DNA sequencer (proviced from Amersham Pharmacia Biotech Ltd.) (Proc. Natl. Acad. Sci. USA 74, 5463-5467, 1977). The 5' upstream region of the human MP52 gene consisting of 3521 bases is presented in SEQ ID No.: 1 of the Sequence Listing. Sequencing was done at least three times.

A site difficult to sequence was subcloned to sequence starting from both terminals. The base sequence from 2880 to 3521 shown in SEQ ID No.: 1 is already reported (Biochem. Biophys. Res. Comm. 204, 646-652, 1994).

5 Example 3: Construction of a recombinant vector containing the human MP52 gene promoter and a luciferase reporter gene

The restriction enzyme map of the 5' upstream region 3.5 kb of human MP52 gene is shown in Fig. 1. An ApaLI-BglI fragment (0.6 kb) of the 5' upstream region of human MP52 gene was purified and the ApaLI end was blunted. Then, this fragment was treated with HindIII and BglII and inserted into a pGL3-basic vector of which HindIII end was blunted. The vector obtained was treated with BglII and KpnI restriction enzymes to ligate to the BglII-KpnI fragment (derived from pBluescript vector) (2.7 kb) of the 5' upstream region of the human MP52 gene. The recombinant expression vector pMP52L (8085 bp) finally obtained contains 3.3 kb of 5' upstream region of human MP52 gene. Fig. 2 shows the constructed recombinant expression vector.

20 Example 4: Measurement of activity of the human MP52 promoter (introduction of the recombinant expression vector into human cells and transient expression)

In order to transiently express the recombinant expression vector pMP52L of human MP52 promoter, said vector 25 was mixed with a vector pRL-SV40 (provided from of Pro Mega Ltd.) containing a sea pansy luciferase gene as an internal control for measurement of introducing efficiency of a gene in an equal quantity. In addition, cationic liposome lipofectamine (provided from Lifetech Oriental Co.) was mixed 30 with said DNA solution to add to human osteosarcoma cells; HOS, MG63, or SaOS-2, for transfection. The fire fly luciferase activity and the sea pansy luciferase activity were measured by Pikka Gene Dual Kit (provided from Toyo Ink The result is shown in Fig. 3. The promoter activity 35 was expressed as a ratio of the fire fly luciferase activity to the sea pansy luciferase activity. From the result, it has been known that the DNA of SEQ ID No.: 1 of the Sequence Listing has a promoter activity.

9

Example 5 : Introduction of the recombinant expression vector into human cells and stabilized expression

In order to express stably the recombinant expression vector pMP52L of human MP52 promoter, said vector was mixed 5 with a plasmid pPUR containing a gene resistant to puromycin in a proportion of 10 : 1, mixed with cationic liposome lipofectamine (provided from Lifetech Oriental Co.) and added to SaOS-2 cells for transfection. The cells into which the target gene has been introduced were selected from a culture medium containing puromycin (provided from Sigma Ltd.).

Example 6 : Screening of active low molecular weight compound

The cells selected were inoculated in a 96-well plate, treated with substances of various chemical compound libraries for 1 - 3 days, dissolved with cytolytic agent (provided from Pro Mega Ltd.), and measured for enzyme activity by employing a luciferase assay kit (provided from Pro Mega Ltd.). By such steps, various substances enhancing or inhibiting the expression of human MP52 can be surveyed.

10

Sequence Listing Free Text

<210> 1

<223> Human MP52 5' upstream gene sequence; the initiating codon ATG is located at the end of the sequence.

<210> 2

<223> Sense PCR primer OD corresponding to human MP52 cDNA sequence

10

<210> 3

<223> Reverse PCR primer OID corresponding to human MP52 cDNA sequence

PCT/IB99/01071 WO 99/66060

11

What is claimed is:

1. A DNA whose nucleotide sequence is represented by the base sequence from No. 1 to No. 3521 in SEQ ID NO.: 1 of the 5 Sequence Listing which encodes a human MP52 gene promoter region, or a fragment thereof.

- 2. A DNA whose nucleotide sequence is represented by the base sequence from No. 1 to No. 2879 in SEQ ID NO.: 1 of the Sequence Listing which encodes a human MP52 gene promoter 10 region, or the fragment thereof.
 - 3. A method for preparing the DNA shown in SEQ ID NO.: 1 of the Sequence Listing comprising the steps of:
 - (1) preparation of a cDNA probe by PCR with a degenerate primer,
- (2) plaque hybridization screening of a human genomic 15 phage library by using the probe, and
 - (3) subcloning of a target fragment from an isolated phage clone.
- A recombinant expression vector characterized by 20 integration of the full length or a part of DNA shown in SEQ ID NO.: 1 of the Sequence Listing into a reporter gene.
 - 5. A method for exploring a useful substance, characterized by using the recombinant expression vector according to claim 4.
- The method for exploring a useful substance according to claim 5, wherein the useful substance is a bone-related substance.
 - The method for exploring a useful substance according to claim 6, wherein the bone-related substance is an
- 30 osteogenesis inducing substance.
 - The method for exploring a useful substance according to claim 6, wherein the bone-related substance is an osteogenesis inhibiting substance.
- The method for exploring a useful substance according to 35 claim 6, wherein the useful substance is a neurogenic substance.
 - 10. The method for exploring a useful substance according to claim 6, wherein the useful substance is an angiogenic

12

substance.

11. The method for exploring a useful substance according to claim 6, wherein the useful substance is an anti-angiogenic substance.

5






FIGURE 1



BEST AVAILABLE COPY



BEST AVAILABLE COPY

FIGURE 3

1

SEQUENCE LISTING

<110> Hoechst Marion Roussel Ltd

<120> Human MP52 gene promoter and method for exploring useful substance by using the same.

<130> JH98K006 PCT SEQUENCES IN ENGLISH

<140>

<141>

<150> 10-170941

<151> 1998-06-18

<160> 3

<170> PatentIn Ver. 2.1

<210> 1

<211> 3521

<212> DNA

<213> HUMAN

<220>

<221> misc_feature

<222> (1)..(3521)

<223> Human MP52 5' upstream gene sequence; the initiating codon ATG is located at the end of the sequence.

2

<400> 1

<400> 1						
tcgactcgat	cactttatgc	aatttaattt	tatttattt	tttagtagag	acagggtttc	60
accatgttag	ccaggatggt	cttgatctcc	tgacctcgtg	atccacctgc	ctcggcctcc	120
caaagtgctg	ggattacagg	cgtgagcact	gcgcctggcc	gcaatttact	tcattgaatc	180
tccaacaaga	gtcctgtgag	gtaagcacta	tcgttatcac	tgtttagagt	ttcagagggg	240
tttagaggct	tcccaagatc	acacaataca	taaatagcag	aaccaagctt	caaaaccagg	300
tctgtttgtc	tccagaatcc	ttgctcttta	tcaagccacg	tacagagaca	ttgtggttgt	360
tcaactcatt	catttattca	ctctctgagc	ttacaaaatg	cttaagaagt	ggcaagacaa	420
ttcttccctt	caagaaactt	agagtctaat	gggaaaggca	ggttatgtcc	acaaataact	480
acacctcaaa	gtagaaaatg	atgatttttg	tcaataaggg	ccagttagag	attaatttat	540
tcatataaca	cgtactgaga	gctgctgtgg	gctgggcctc	tgccaagcac	tgggtataat	600
tcaaagataa	atatggcaca	gtctagtcat	aatctaatgg	gagagacagg	tatgtaaaga	660
aattattata	gttataagga	attaagttat	gattttaaaa	tatgacaaaa	tacagagtgg	720
aagaaacaaa	ctgtgttggg	atagggggca	gaggcagaga	ctgaataaac	tttcagctgc	780
caaaaaatgt	aaaagagcac	agaaaatgtg	tgatggcagc	tcagggggag	tgtgcatttt	840
ctggaggatg	caagaaggct	tcatggagga	ggtggcacgt	gggttggcac	ttgtttgatg	900
ggtagctttt	agtagagggt	tgaaccagag	agagcctgga	ggcctagaag	ccagttatag	960
gggagaatca	tagcgaggca	agcaatgagg	aggacagggc	agtgacagag	gtggtgagga	1020
gggagtgaag	ggagggatgc	cagagaggac	tggttcagtg	ggaacaagtc	tgtttttaat	1080
ttttttattt	tattaatatt	atttatttac	ttattttgtg	tgtgtgtgat	ggactctagc	1140
tctgtcacct	aggctggagt	gcagtggtgc	gatctcggct	cactgcaacc	tttgcctcca	1200
cacccggcta	attttttgta	tttttggtag	agatgcctgc	aatcccagca	ctttgagagg	1260
ccaaggtggg	tggatcactt	gaggtcagga	gctcgagacc	agactggcca	acatggtgaa	1320
actccgtctc	tactaaaaat	acaaaaaata	gataggcatg	gtggtgtgca	cctgtagtcc	1380
cagctactcg	agaggctgag	gcaggagaat	cgtttgaacc	caggaggtgg	agattacagt	1440
gagccaagat	cgcaccactg	cactccagcc	taggtgacag	agtgagactc	tgtctcaaaa	1500
aaaaaaaaa	aaaaaaaaa	aaagaaggaa	gcctatctga	ctattgcttc	tcccccgcca	1560
tccttcttac	agcgtgaaaa	gtgttgttgt	agagaaatgt	acagaagggg	taactggctc	1620
tgtctgggga	cgaggaaagg	cttcccgaag	gaggtaaatt	ttcagctaaa	cttctcagga	1680
tgtagagtgc	ttcctccagt	tgtgggagag	agagaggagg	atatttcagg	tgaagcgaac	1740
agctttcccc	gctgaccctg	tgttaagttg	cccaggggct	ctacggcgaa	ggttccccag	1800
aggaagggat	cttccatact	tcaatgaggc	actttaagaa	acccttgagt	tcagctggaa	1860
ttcagacctt	ccagagctac	cagaaaaaca	tcatgtggga	aattgtgccc	ggcgcggtgt	1920
cacgcctgta	atcccagcac	tttgggaggc	cgaggcgggc	ggatcacgag	gtcaggagat	1980
cgagaccaac	ctggctaaca	tggtgaaacc	ccatctctac	tacaaacgta	aaaaatcagc	2040
caggtgtggt	ggcaggggcc	tgtagtccca	gctactgggg	aggctgaggc	aggggaatgg	2100
cgtgaacctg	ggaggcagag	attgcagtga	gctgagatca	tgccactgca	ctcgacagag	2160
gcgaactccg	tctcaaaaaa	aaaaaaaaa	aaagcgtcat	gtgggaaatg	gcttttccag	2220

3

```
cctcctgtag gggccgctgc tgccccagac tcagccagtc ttgtctaaga aaactcagag 2280
gacgtctctg ctggggtggt ggtggggtac accctggacc tgccgccatc caaggagtga 2340
acttaagggc gacagtgccc ccaagcctga agaatatgca ctcagtcaat ggcatttggg 2400
ggtagggagg gggtgtagtg ggcactgata tttttcagtc tctgggtcac cacaagttta 2460
ttaaaataaa agaaaatgtg ggtaaatgct gccatctgtg ctgtccctac tcccaataca 2520
cacacacaca cacacacaca cacacacaca cacacacacg cataacaaat caggtcccag 2640
actagatgca ggagttgaaa ctgctttaaa acactctaag aacttaacta caaacccaca 2700
ctccccatgc ttctggggct taagatcttt gagctcattc ttcagcatct ctccacgaga 2760
aagtggggtg ggctgctcca tgaggtggag gtgaagaccc ctgagtctgc cccgtggagg 2820
gggaggeece ctaageetag agteeegetg cagggetetg tgeeaggage ceeegtgage 2880
catggcctcg aaagggcagc ggtgattttt ttcacataaa tatatcgcac ttaaatgagt 2940
ttagacagca tgacatcaga gagtaattaa attggtttgg gttggaattc cgtttccaat 3000
tectgagtte aggtttgtaa aagattttte tgageacetg caggeetgtg agtgtgtgtg 3060
tgtgtgtgtg tgtgtgtgt tgtgtgtgaa gtattttcac tggaaaggat tcaaaactag 3120
ggggaaaaaa aaactggagc acacaggcag cattacgcca ttcttccttc ttggaaaaat 3180
ccctcagcct tatacaagcc tccttcaagc cctcagtcag ttgtgcagga gaaagggggc 3240
ggttggcttt ctcctttcaa gaacgagtta ttttcagctg ctgactggag acggtgcacg 3300
tetggatacg agageattte cactatggga etggatacaa acacacacce ggeagaette 3360
aagagtetea gaetgaggag aaageettte ettetgetge tactgetget geegetgett 3420
ttgaaagtcc actcctttca tggtttttcc tgccaaacca qaggcacctt tgctgctgcc 3480
gctgttctct ttggtgtcat tcagcggctg gccagaggat g
                                                               3521
```

<210> 2

<211> 36

<212> DNA

<213> HUMAN

<220>

<221> misc_feature

<222> (1)..(36)

<223> Sense PCR primer OD corresponding to human MP52 cDNA sequence

<400> 2

atgaattccc atggacctgg gctggmakga mtggat

4

<211> 29

<212> DNA

<213> HUMAN

<220>

<221> misc_feature

<222> (1)..(29)

<223> Reverse PCR primer OID corresponding to human MP52 cDNA sequence

<400> 3

atgaattcga gctgcgysgg srcacagca

29

INTERNATIONAL SEARCH REPORT

Internat | Application No PCT/IB 99/01071

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/85 C120 C12Q1/68 According to International Patent Classification (IPC) or to both national classification and IPC B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C12N C12Q IPC 6 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) C. DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. χ HOTTEN G ET AL: "Cloning and expression 1 of recombinant human growth/differentiation factor 5" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS., vol. 204, no. 2, 28 October 1994 (1994-10-28), pages 646-652, XP002115693 cited in the application figure 1 2-11 WO 95 04819 A (BIOPH BIOTECH ENTW PHARM X 1 GMBH) 16 February 1995 (1995-02-16) Υ Seq.Id.No.1 2-11 -/--Further documents are listed in the continuation of box C. Patent family members are listed in annex. Special categories of cited documents: "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance invention "E" earlier document but published on or after the international "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such docucitation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means ments, such combination being obvious to a person skilled in the art. "P" document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 27 September 1999 08/10/1999 Name and mailing address of the ISA Authorized officer European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo ni, Lonnoy, 0 Fax: (+31-70) 340-3016

INTERNATIONAL SEARCH REPORT

Interna: I Application No
PCT/IB 99/01071

212		PCT/IB 99/01071
Category *	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	
Calegory	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Х	WO 97 03188 A (BIOPH BIOTECH ENTW PHARM GMBH ;HOETTEN GERTRUD (DE); POHL JENS (DE) 30 January 1997 (1997-01-30)	1
Y	Seq.Id.No.2	2–11
X	WO 97 04095 A (MATSUMOTO TOMOAKI ;HOECHST JAPAN (JP); KIMURA MICHIO (JP); FUJINO) 6 February 1997 (1997-02-06)	1
Y .	Seq.Id.No.1	2-11
Y	SAMBROOK J ET AL: "MOLECULAR CLONING: A LABORATORY MANUAL, PASSAGE" MOLECULAR CLONING,1987, page 2.114-2.117; 14.7-14.8 XP002115694 the whole document	2,3
Y	WO 98 23740 A (AKZO NOBEL NV ;DIJKEMA REIN (NL); DEN WIJNGAARD ARTHUR VAN (NL)) 4 June 1998 (1998-06-04) claim 5	4-11
Υ .	WO 97 15308 A (ZYMOGENETICS INC; OSTEOSCREEN INC (US); UNIV TEXAS AT AUSTIN (US)) 1 May 1997 (1997-05-01) cited in the application page 5, paragraph 2; examples 2,4	4-11
	,	
		·

INTERNATIONAL SEARCH REPORT

is....mation on patent family members

Internal | Application No | PCT/IB 99/01071

				99/010/1	
Patent document cited in search report		Publication date		atent family member(s)	Publication date
WO 9504819	Α	16-02-1995	DE	4420157 A	23-02-1995
			AÜ	688362 B	12-03-1998
			AU	7498694 A	28-02-1995
			CA	2169171 A	16-02-1995
			CN	1129013 A	14-08-1996
			CZ	9600357 A	17-07-1995
			EP	0713529 A	29-05-1996
			HÚ	74271 A	28-11-1996
			JP	9501053 T	04-02-1997
•			NZ	271376 A	24-04-1997
			ZA	9405992 A	14-03-1995
				3403332 K	14-03-1995
WO 9703188	Α	30-01-1997	DE	19525416 A	16-01-1997
	••	00 02 2007	AU	6615196 A	10-02-1997
			EP	0837938 A	29-04-1998
		•	JΡ	11509097 T	17-08-1999
			·	~~~~~~	17 00 1999
WO 9704095 ·	Α	06-02-1997	JP	9031098 A	04-02-1997
			AU	704364 B	22-04-1999
			AU	6530496 A	18-02-1997
			CA	2224289 A	06-02-1997
			CN	1196087 A	14-10-1998
			EP	0866125 A	23-09-1998
			HU	9901204 A	28-07-1999
			NO	980300 A	23-01-1998
		~~~~~~~			
WO 9823740	Α	04-06-1998	AU	5558498 A	22-06-1998
			.EP	0937145 A	25-08-1999
WO 9715308	Α	01-05-1997	AU	706262 B	10-06-1999
3/13300	Λ	01 03 133/	AU	7471096 A	15-05-1997
			CA	2235481 A	01-05-1997
			CN	1201393 A	01-05-1997
			CZ	9801153 A	
			EP		16-12-1998
		•		0866710 A	30-09-1998
			HU	9802319 A	01-02-1999
			NO PL	981810 A 327617 A	22-06-1998
			r:	3//DI/ A	21-12-1998